

Impacts of a Workplace-Based Weight-Control Intervention on Objective and Perceived Physical Activity among a Subgroup of Workers

Rachel Tabak¹, Ryan Colvin², Jaime R. Strickland², Ann Marie Dale², Maura Kepper¹, Tara Ruggeri²,
and Bradley Evanoff²

¹Brown School at Washington University in St. Louis, St. Louis, MO

²Washington University School of Medicine in St. Louis, St. Louis, MO

Abstract

Physical activity (PA) has many benefits; however, groups facing barriers to health-promoting behaviors are less likely to be physically active. This may be addressed through workplace interventions. The current study employs objective (accelerometry) and perceived (International Physical Activity Questionnaire [IPAQ]) measures of PA among a subset of participants from the “Working for You” study, which tests a multi-level (work group and individual) workplace intervention targeted at workers with low-incomes. Linear mixed and hierarchical logistic regression models are used to determine the intervention’s impact on moderate- to vigorous-PA (MVPA) and achieving the PA Guideline for Americans (≥ 150 minutes MVPA/week), respectively from baseline to 6- and 24-months, relative to a control group. Correlations (Spearman Rho) between perceived and objective PA are assessed. Of the 140 workers (69 control, 71 intervention) in the sub-study, 131 (94%) have valid data at baseline, 88 (63%) at 6-months, and 77 (55%) at 24-months. Changes in MVPA are not significantly different among intervention relative to control participants assessed by accelerometer or IPAQ at 6- or 24-months follow-up. The percent achieving the PA Guideline for Americans does not vary by treatment group by any measure at any time point (e.g., baseline accelerometry: [control: n=37 (57%); intervention: n=35 (53%)]). This study identifies limited agreement (correlation range: 0.04 to 0.42, all $p > .05$) between perceived and objective measures. Results suggest the intervention did not improve PA among the sub-study participants. Though agreement between objective and perceived MVPA is low, similar conclusions regarding intervention effectiveness are drawn.

Trial registration: ClinicalTrials.gov: NCT02934113

Keywords: physical activity, measurement, worksite

Physical activity (PA) promotes health, prevents obesity, and reduces the risk of diabetes and other chronic conditions (Piercy et al., 2018; United States Department of Health and Human Services, 2020). The 2019 Physical Activity Guidelines for Americans identified numerous benefits of being active including: prevention of numerous chronic diseases (e.g., cancer; heart disease, type 2 diabetes), excessive weight gain, and depression (United States Department of Health and Human Services, 2020). Unfortunately, like many health behaviors that can help prevent chronic disease, individuals from historically marginalized groups face barriers (e.g. discretionary time, physical safety) to health and health promoting behaviors and are less likely to be physically active than those with fewer barriers (Haskell et al., 2007; Troiano et al., 2008; Tucker et al., 2011; Williams et al., 2018). This contributes to a disparate chronic disease burden (Barbeau et al., 2004; Byers et al., 2008; Clegg et al., 2009; Goetzl & Ozminkowski, 2008; Huang et al., 2011; Kogevinas & Porta, 1997; McLaren, 2007; Woods et al., 2006). These barriers are disproportionately experienced by populations that have been historically marginalized (e.g., racial/ethnic minorities) and contribute to chronic disease disparities (Hawes et al., 2019; Howell & Booth, 2022; Z. Javed et al., 2022; Javed et al., 2022; Luo et al., 2022; Prener et al., 2021).

Interventions are needed to promote PA among populations who face great barriers and who are less likely to have access to and benefit from health systems or community-based PA interventions (Lee & Cubbin, 2009; Mazzucca et al., 2021). Non-clinical settings such as workplaces offer a potentially effective setting for health promotion programs to reach and impact a broader, more diverse population (Mazzucca et al., 2021; Parrish et al., 2018). However, studies testing healthy weight interventions often have limited racial/ethnic representation (Aneni et al., 2014; Brown et al., 2018; Coenen et al., 2020; Haughton et al., 2018; Rongen et al., 2013). It is particularly important that interventions be developed in workplaces with employees who are low-income and who disproportionately face barriers to health behaviors and have previously had more limited access to workplace interventions (Grosch et al., 1998; Kopicki et al., 2009; Linnan et al., 2001; Parrish et al., 2018). There is mixed evidence regarding the effectiveness of workplace interventions to change health behaviors and address inequities, suggesting further work in this area is needed (Cairns et al., 2015; Coenen et al., 2020).

The “Working for You” study aimed to test a workplace intervention targeted at workers who were low-income in partnership with an academic medical center. Working for You adapted existing group- and individual-level intervention models to promote healthy weight and prevent chronic disease among workers (Stein et al., 2019; Tabak et al., 2020; Tabak et al., 2018). Though the primary target of this multi-level intervention was weight, an important behavioral component was increasing moderate- to

vigorous-PA (MVPA). Given the robust discussion in the literature regarding appropriate methods to assess MVPA the current study employed both objective (accelerometry) and perceived (International Physical Activity Questionnaire [IPAQ]) measures of PA among a subset of participants from the larger multi-level trial to a) determine the impact of the intervention on MVPA and b) assess the agreement between these measures in this working population (Dowd et al., 2018; Kelly et al., 2016; Limb et al., 2019),.

Methods

Participants and Intervention

Participants for this study were recruited from a large cluster-randomized trial testing the impact of Working for You, a multi-level weight loss intervention among approximately 1,000 employees across 22 work groups at a Midwestern academic medical center. The study was cluster-randomized at the work group level, specifically targeting work groups where the average work group salary was less than \$50,000 per year. Participants were predominantly employed in office and administrative support (e.g., billing and account collectors, medical claims), medical and lab technicians, housekeeping, and food service. The trial and intervention have been described elsewhere (Stein et al., 2019; Tabak et al., 2020; Tabak et al., 2018). Briefly, all participants in work groups randomized to the intervention received the group-level component, which is a workplace participatory program involving worker teams. These teams engaged in the design and implementation of weight-reducing interventions through changes in their workplace environments. Each work group developed their own intervention, with considerable variation (e.g., walking club, healthy snack provision, distribution of more information on employer wellness resources). Employees in the work groups randomized to intervention with obesity (body mass index ≥ 30) were also invited to participate in an interactive obesity treatment approach (iOTA). iOTA includes a) quarterly health coach meetings during which the participant collaboratively sets goals and b) an automated interactive SMS text-messaging program for ongoing support and self-monitoring (Stein et al., 2019; Tabak et al., 2018). Participants in work groups randomized to the control group completed data collection and did not have work group level or iOTA interventions. The larger study excluded those with a current cancer diagnosis or treatment, pregnant at study recruitment, or who had bariatric procedure within the last year (Stein et al., 2019). The current sub-study recruited participants from the larger study (as determined by a power calculation) with obesity (body mass index ≥ 30) and with no plans to leave the work group within the next six months.

While the Working For You study did not specifically target MVPA as an outcome, assessing MVPA across treatment groups was a reasonable objective for this sub-study. Eighty-five percent of sub-study participants in the intervention group were also enrolled in iOTA. iOTA

participants could set goals based on a variety of weight loss behaviors. Goals targeting physical activity (e.g., increase number of steps per day, increase brisk activity) were chosen by 88% of participants over the course of the study, with the proportion being 100% among those contributing data to the final follow up time point.

Measurement Procedures

Individual meetings were scheduled with each recruited participant, which included an informed consent process and explanation of the accelerometer device, the device wear log, and an instruction sheet. A member of the research team also showed each participant how to properly wear the accelerometer. Participants completed study assessments for the current sub-study at three time-points: baseline (enrollment), 6-months, and 24-months; however, if a participant did not have valid data at the baseline and 6-month time-points, they were not asked to participate in the 24-month time-point. Data was collected between April 2017 and December 2019.

Accelerometry

At baseline, 6-, and 24-months, free-living physical activity was measured with an ActiGraph GT3+ accelerometer (Ft. Walton Beach, FL) that was belt worn on the right hip during waking hours. Participants were instructed to wear the device for \geq seven days (with \geq three of the seven days being days they were at work) and to remove the device for showering or other water activities (Troiano et al., 2008). Participants were asked to use the device log to record the days and times they wore the accelerometer. Device log data was used during data reduction in order to visualize what days/time, and how much time, each participant spent wearing the devices at work, not at work, and total during the duration of the study.

Survey Data

Participants' demographic characteristics were drawn from their baseline survey (a self-report instrument) for the larger trial. At the end of each data collection time-point (baseline, 6-months, and 24-month follow up), the short version of the International Physical Activity Questionnaire (IPAQ) was self-administered to each participant (<https://sites.google.com/site/theipaq/>). The IPAQ is a self-reported questionnaire that measures perceived moderate and vigorous PA and sedentary behavior (Craig et al., 2003). Participants completed the IPAQ which used a referent period of the previous week (corresponding to the time they wore the accelerometer). Daily estimates of MVPA (weekly estimates divided by seven) were derived to put IPAQ data into same units as the actigraphy data. Participants who reported over 720 minutes (12 hours) per day were excluded from analyses using IPAQ data.

Analysis

ActiGraph data were downloaded and scored using the ActiLife v6.7.3 software. Data decision rules regarding

non-wear time (sequences of \geq 90 min of 0 counts) and the definition of a valid day (\geq 10 hours of waking-wear time in a 24-hour period) were used (Troiano et al., 2008). To determine waking-wear time, data points were excluded before and after participant reported time they applied/removed the accelerometry device. Participant data were included in the analysis if they have a minimum of five days of recorded activity with an average \geq 600 minutes of recorded activity per day or for four days with \geq 3000 total minutes (James et al., 2017). Participants wore the devices for 784 ± 76 min/day over 6.8 ± 1.0 days/week. Intensity cut points were calculated using the Troiano standard adult thresholds, which identified sedentary activity as 0-99 CPM (counts per minute), light activity as 100-2019 CPM, moderate activity as 2020-5998 CPM, and vigorous activity as \geq 5999 CPM (NCI Division of Cancer Control and Population Sciences: Epidemiology and Genomics Research Program, 2020). Moderate to vigorous bouts were defined as having a minimum of 10 minutes of activity with a bout of 2 minutes and a minimum of 2,020 counts per minute (Matthew, 2005; Matthews et al., 2008; Troiano et al., 2008; Tudor-Locke et al., 2010).

Differences in baseline characteristics were assessed using Fisher's exact test and independent sample T-tests. Change in MVPA over time was assessed for two outcomes, each assessed using linear mixed models; objectively-measured (using accelerometry) and perceived activity (using the IPAQ). These models included MVPA (based on either actigraphy or IPAQ) as the dependent variable and treatment group, time point (baseline, 6-months, and 24-months) and the interaction of treatment group and time point as the independent variables (fixed effects). The accelerometry-based data analysis also included an additional fixed effect for average daily wear time, to account for varying wear times across individuals. Random intercepts for work group were included to account for the hierarchical structure of the data.

The proportion of participants meeting US Department of Health and Human Services Physical Activity Guideline for Americans, 2nd edition guidelines for MVPA were calculated by dichotomizing objectively measured and perceived MVPA at minutes/week \geq 150 (Piercy et al., 2018). Differences in proportion of participants meeting guidelines by treatment group at each time point were assessed using logistic regression models, where guideline status (met vs not met) was the dependent variable, and treatment group, time point, and the interaction of treatment group and time point were the independent variables.

In all models (linear and logistic) the hierarchical study design was accounted for using random intercepts for work group, and a first-order autoregressive covariance structure was applied to the residuals to account for repeated measures of the outcome among participants (Singer et al., 2003). This group-level analysis incorporates group mean comparisons, rather than individual trajectories of analysis, and allows for inclusion of participants providing valid data at selected time-points. Adjusted models also included additional fixed effect covariates age, annual income, and


baseline body mass index, as these were not balanced by randomization. Finally, Spearman Rho correlations were used to assess agreement between objectively measured and perceived activity for MVPA. Agreement was assessed at each time point (baseline, 6-months, and 24-months), overall and stratified by treatment group. All analyses were conducted using SAS Software 9.4 (The SAS institute, Cary, NC).

Results

Of the 237 participants determined to be eligible for this study, 176 provided informed consent, and among these, 140 had accelerometer data that met wear-time criteria (Figure 1). Of these, 131 (94%) had valid baseline data, 88 (63%) had valid data at 6-months and 77 (55%) had valid data at 24-months. There was a statistically significant difference in follow-up between groups at 24 months, with greater participation among control participants; however, the difference was not statistically significant at 6 months or “any follow up” (6 or 24 months).

Figure 1

Participant flow in the subgroup of Working for You study eligible for the accelerometry sub-study



Demographic characteristics at baseline for the 140 participants included in the analysis are presented in Table 1. Participant age differed between groups, with the mean age in the control group 46.1 ± 10.7 years and 41.5 ± 10.1 years in the intervention group. The average body mass index for participants was greater (39.0 ± 7.2) for those in the intervention group compared to the control group (36.0 ± 5.0 , $p=0.005$). Most participants identified as Black or African American ($n = 71$, 51%) or White or Caucasian (n

$= 64$, 46%). The intervention and control groups differed with regard to education; while a similar percent (15% in both groups) reported a high school education or lower, 56% of those in the control group and 75% in the intervention group reported technical school or some college, and college/post-graduates were represented 29% of the control group, but only 10% of the intervention group. Across the groups, most participants (64%) reported a household income less than \$70,000 per year.

Table 1. Baseline characteristics for participants in the sub-group of the Working For You study

	Total N = 140*	Control N = 69	Intervention N = 71	P†
Weight, kg	103.7 ± 17.8	100.6 ± 15.7	106.8 ± 19.3	0.09
BMI	37.5 ± 6.3	36.0 ± 5.0	39.0 ± 7.2	0.005
Female gender	102 (73.4)	48 (70.6)	54 (76.1)	0.57
Age, years	43.7 ± 10.6	46.1 ± 10.7	41.5 ± 10.1	0.009
Race				0.37
Black/African American	71 (50.7)	33 (47.8)	38 (53.5)	
White/Caucasian	64 (45.7)	32 (46.4)	32 (45.1)	
Other	5 (3.5)	4 (5.8)	1 (1.4)	
Hispanic	1 (0.7)	0 (0.0)	1 (1.4)	1.00
Education				0.01
High school or lower	20 (14.6)	10 (14.7)	10 (14.5)	
Technical school/some college	90 (65.7)	38 (55.9)	52 (75.4)	
College graduate or post-graduate	27 (19.7)	20 (29.4)	7 (10.1)	
Annual household income				0.17
≤\$30,000	26 (19.0)	10 (14.9)	16 (22.9)	
\$30,001-\$70,000	62 (45.3)	28 (41.8)	34 (48.6)	
≥\$70,001	49 (35.8)	29 (43.3)	20 (28.6)	
Average daily accelerometer wear time, minutes	772.5 (729.1, 817.6)	756.9 (724.7, 809.3)	784.9 (734.3, 825.0)	0.18

Note. Presented are N (%) for categorical variables and mean ± SD or Median (25th, 75th %tile) for continuous variables

*Nine participants did not provide valid accelerometry data at baseline, but did contribute at later time points, so were included in this study.

†based on Fisher's exact test (categorical variables), two sample T test or Wilcoxon rank sum test (continuous variables), bold indicates p<.05

There were no statistically significant differences between groups in MVPA at baseline assessed by accelerometry (control group mean [95% confidence interval]: 30.1 [22.9, 37.3]); (intervention group 25.9 [18.4, 33.5], Table 2). From baseline to 6-months, objective MVPA decreased by 1.8 minutes (95% confidence interval: -7.2, 3.5) for the control group, but only 0.2 minutes (-5.9, 5.6) for the intervention group and from baseline to 24-

months MVPA decreased by 4.2 minutes (-8.5, 0.1) control group, but only 0.4 minutes (-5.6, 4.9) for the intervention group. When the difference in change from baseline between groups (i.e., intervention relative to control) was estimated using a regression model with age and income, the difference was not statistically significant at 6-months (1.7 minutes [-6.2, 9.5]) or at 24-months (3.8 [-2.9, 10.6]).

Table 2. Change in objective (accelerometry) and perceived (International Physical Activity Questionnaire, IPAQ) moderate or vigorous physical activity (MVPA) by treatment group

	MVPA (average minutes/day) mean (95% CI)*	Difference between treatment groups, Intervention vs Control (95% CI)	Unadjusted P†	Adjusted P‡
Objective MVPA- Accelerometry, controlling for wear time				
<i>Baseline</i>				
Control	30.1 (22.9, 37.3)	<i>ref.</i>		
Intervention	25.9 (18.4, 33.5)	-4.2 (-14.6, 6.3)	0.41	0.20
<i>Change from baseline to 6-month follow up</i>				
Control	-1.8 (-7.2, 3.5)	<i>ref.</i>		
Intervention	-0.2 (-5.9, 5.6)	1.7 (-6.2, 9.5)	0.68	0.77
<i>Change from baseline to 24-month follow up</i>				
Control	-4.2 (-8.5, 0.1)	<i>ref.</i>		
Intervention	-0.4 (-5.6, 4.9)	3.8 (-2.9, 10.6)	0.27	0.36
Perceived MVPA - IPAQ				
<i>Baseline</i>				
Control	126.1 (63.7, 188.5)	<i>ref.</i>		
Intervention	132.0 (66.1, 198.0)	5.9 (-84.9, 96.7)	0.89	0.73
<i>Change from baseline to 6-month follow up</i>				
Control	-10.1 (-59.5, 39.4)	<i>ref.</i>		
Intervention	57.4 (6.2, 108.5)	67.4 (-3.7, 138.6)	0.06	0.12
<i>Change from baseline to 24-month follow up</i>				
Control	-27.0 (-70.8, 16.9)	<i>ref.</i>		
Intervention	10.8 (-42.0, 63.6)	37.7 (-30.9, 106.4)	0.28	0.51

Note. *Mean changes and CIs are estimated from the regression model. They take hierarchical structure into account and will vary slightly from raw means.

†P values derived from linear mixed model with MVPA as the dependent variable and treatment group, time point (baseline, 6, 24 months) and the interaction of treatment group and time point the independent variables (fixed effects). Random intercepts for work group were included to account for the hierarchical structure of the data. Repeated MVPA measures over time were accounted for using first order autoregressive covariance structure applied to the residuals.

‡P values based on model with structure in unadjusted models, plus additional fixed effect covariates age, annual income and baseline body mass index.

Ns = 140 participants contributed accelerometry data at 293 time points (baseline, 6-, 24-months); 139 participants contributed IPAQ data at 288 time points (baseline, 6-, 24-months)

The trends were similar using perceived (IPAQ) measure, however the number of minutes as well as the confidence intervals were much larger: difference in change from baseline between groups estimated using a regression model (67.4 [-3.7, 138.6]) at 6-months and (37.7 [-30.9, 106.4]) at 24-months.

The baseline distribution of those meeting the Physical Activity Guideline for Americans did not vary by treatment

group based on accelerometry: 37 (57%) for control group; 35 (53%) for intervention group (Table 3). Similar results were found with perceived activity, although the proportion reporting meeting guidelines was higher than was identified by accelerometry (47 [73%] for control group; 44 [70%] for intervention group). The proportion meeting guideline recommendations did not vary by treatment group at the follow up time points, when assessed by accelerometry or perceived (IPAQ).

Table 3. Proportion of participants meeting MVPA guidelines* by treatment group and time point as measured by objective accelerometry and perceived International Physical Activity Questionnaire (IPAQ)

	Proportion meeting guidelines*, N (%)	OR (95% CI)*	Unadjusted P*	Adjusted P†
Objective MVPA- Accelerometry				
<i>Baseline</i>				
Control	37 (56.9)	<i>ref.</i>		
Intervention	35 (53.0)	1.10 (0.28, 4.34)	0.88	0.75
<i>6 month follow up</i>				
Treatment group				
Control	26 (55.3)	<i>ref.</i>		
Intervention	26 (63.4)	1.72 (0.40, 7.48)	0.45	0.92
<i>24 month follow up</i>				
Control	22 (45.8)	<i>ref.</i>		
Intervention	11 (37.9)	0.89 (0.20, 4.01)	0.87	0.32
Perceived MVPA - IPAQ				
<i>Baseline</i>				
Control	47 (73.4)	<i>ref.</i>		
Intervention	44 (69.8)	0.85 (0.21, 3.38)	0.80	0.92
<i>6 month follow up</i>				
Control	31 (70.5)	<i>ref.</i>		
Intervention	33 (80.5)	1.58 (0.35, 7.11)	0.54	0.51
<i>24 month follow up</i>				
Control	31 (66.0)	<i>ref.</i>		
Intervention	24 (82.8)	2.62 (0.57, 12.04)	0.20	0.20

Note. *Relative Odds and P values derived from generalized linear mixed model (logistic regression) with meeting moderate to vigorous physical activity (MVPA) guidelines (≥ 150 min MVPA/week) as the dependent variable and treatment group, time point (baseline, 6-, and 24-months) and the interaction of treatment group and time point the independent variables (fixed effects). Random intercepts for work group were included to account for the hierarchical structure of the data. Repeated MVPA measures over time were accounted for using a first order autoregressive covariance structure applied to the residuals.

†P values based on model with structure in unadjusted models, plus additional fixed effect covariates age, annual income and baseline body mass index. .

Ns = 140 participants contributed accelerometry data at 296 time points (baseline, 6-, and 24-months); 139 participants contributed IPAQ data at 288 time points (baseline, 6-, and 24-months)

When exploring the correlations between accelerometer- and perceived-MVPA, the correlations were low (range: 0.04 for the control group at 24-months to 0.42 for

intervention group at 24-months), with no statistically significant correlations (i.e., all adjusted and unadjusted $p > .05$, Table 4).

Table 4. Correlation of Objective (Accelerometry) and Perceived (International Physical Activity Questionnaire, IPAQ) moderate or vigorous physical activity (MVPA) by treatment group

	Spearman Rho	Unadjusted P*	Adjusted P†
<i>Baseline</i>			
Overall	0.18	0.59	0.51
By treatment group			
Control	0.16	0.64	0.69
Intervention	0.22	0.76	0.60
<i>6 month follow up</i>			

Overall	0.28	0.76	0.72
By treatment group			
Control	0.29	0.79	0.85
Intervention	0.29	0.88	0.73
<i>24 month follow up</i>			
Overall	0.14	0.79	0.61
By treatment group			
Control	0.04	0.64	0.57
Intervention	0.42	0.35	0.22

Note. *P values derived from linear mixed model with objective (accelerometry) MVPA as the dependent variable and treatment group, time point (baseline, 6-, and 24-months), perceived (IPAQ)MVPA, interactions treatment group by time point, treatment group by IPAQ-measured MVPA, time point by IPAQ-measured MVPA, and treatment group by time point by perceived (IPAQ) MVPA as independent variables (fixed effects). Random intercepts for work group were included to account for the hierarchical structure of the data. Repeated MVPA measures over time were accounted for using a first order autoregressive covariance structure applied to the residuals.

†P values based on model with structure in unadjusted models, plus additional fixed effect covariates age, annual income and body mass index.

Ns = 139 participants contributed accelerometry data at 288 time points (baseline, 6-, and 24-months)

All results were similar when intervention group participants were limited to those in iOTA (data not shown), with 82% of intervention group participants also in iOTA.

Discussion

This study explored the impact of a multi-level weight loss intervention, Working for You, on PA, among a subgroup of participants, who wore accelerometers. The analysis did not detect significant differences in changes in MVPA among intervention group participants relative to those in the control group assessed by both accelerometry and IPAQ at 6- or 24-months follow-up. Further, though similar intervention affects were concluded based on these two measurement methods, the study identified limited agreement between the objective (accelerometry) and perceived (IPAQ) assessment of MVPA.

Working for You was developed to include many evidence-based strategies (e.g., tailored behavioral goals, skills training, informational and motivational text messages, behavior self-monitoring delivered by SMS text-messaging, and workplace participatory program) to promote health behavior changes, including increase in PA, related to weight loss and prevention of chronic disease (Stein et al., 2019; Strickland et al., 2020; Tabak et al., 2018). The null findings are consistent with the literature, which shows limited effectiveness of work-place health promotion interventions on PA behavior (Malik et al., 2014), particularly those delivered through technology (Aneni et al., 2014; Direito et al., 2017) and those targeting employees working in jobs that pay lower wages (Coenen et al., 2020; Rongen et al., 2013) and those with low socioeconomic status (even when they are effective in those with higher socioeconomic status) (Western et al., 2021). In the current study, the null findings may be related to the

low intensity intervention overall, and a broad focus on (and powered to detect) weight loss, rather than on PA specifically. While participants were encouraged to select a PA goal as one of the three goals they aimed to work toward, not all participants chose this focus (only 12% of participants never selected a PA goal). In addition, while participants engaged with the individual-level iOTA intervention, preliminary findings suggest there were relatively few changes implemented in the workplace environment resulting from the work group-level intervention, which may contribute to limited impact on individual behavior. Interventions, which comprehensively change the workplace environment (and likely broader environmental and policy conditions) may be needed in addition to more intensive interventions targeting individual drivers of behaviors (Gu et al., 2020; Song & Baicker, 2019; Sorensen et al., 2016; Terry, 2019; Zhu et al., 2020).

Future research in PA promotion among diverse communities can consider the findings of the current study. The study did not demonstrate an impact on PA, which is notable given participants were more diverse with respect to numerous characteristics (e.g., race, income, education) than those in most worksite PA studies, including those in medical settings (Aneni et al., 2014; Blake et al., 2017; Brown et al., 2018; Coenen et al., 2020; Haughton et al., 2018; Rongen et al., 2013). Numerous structural factors (e.g., time, safe places to be active, social networks) influence, and disproportionately hinder, activity among individuals impacted by disparities (Barbeau et al., 2004; Huang et al., 2011; McLaren, 2007). More intensive or alternative strategies may be needed to overcome such structural factors and other barriers both inside and outside the workplace (Lee & Cubbin, 2009). Data were not collected on the participants' home and neighborhood environments, which limits the understanding of broader

potential impacts on PA behaviors. However, there is ample evidence of the importance of neighborhood environments (Kepper et al., 2019; Smith et al., 2017) and activity and the importance of place-based interventions to reduce health inequities (McGowan et al., 2021). In addition, lower income workers are more likely to be facing barriers outside the workplace (e.g., financial, environmental or social stressors) which can take precedence over preventive behaviors (Kreuter et al., 2016; Thompson et al., 2019).

When considering the null findings in the current study, it is important to consider the high level of PA among this group of participants; at baseline 55% engaged in MVPA (assessed by accelerometer) for at least 150 minutes per week (achieving the MVPA recommended by the 2018 PA guidelines) (Piercy et al., 2018). These high levels of PA are likely due to physical work demands, including work within a large physical campus. An intervention study by Folta et al. found approximately 20% of participants engaged in 150 minutes of MVPA per week (2019). However, a study of hospital workers found 62 out of 63 participants (98%), 21 of 22 (96%) physicians, and 100% of nurses and supporting staff, spent over 300 min in MVPA per week (Jun et al., 2019). These already high levels of MVPA may make additional activity challenging to achieve.

This study is strengthened by the randomized design with an intent-to-treat analysis, and the assessment of PA using both objective and perceived measures. Though agreement between objective and perceived activity was low, the current study drew similar conclusions regarding intervention impact with both accelerometry and IPAQ measures, which was consistent with a recent analysis of the Diabetes Prevention Follow-up or Outcomes Study (Kriska et al., 2020). This contributes to the discussion of whether perceived activity data, rather than requiring resource and time intensive objective assessments may be useful in intervention studies. However, other intervention studies comparing objective and perceived activity, with larger sample sizes have found that this may not always be the case (Limb et al., 2019). An important limitation of the current study is the inclusion of only a subset of the total participants in the cluster randomized study, limiting the sample size and increasing the potential for selection bias, and the significant attrition throughout the 24-month follow-up. These limit both power and generalizability. As the primary study was not powered to detect an effect in physical activity in a sub-set of participants and there was a lack of balance achieved by randomization between demographic characteristics in the control and intervention groups in the pilot sample, it is possible these are reasons an intervention effect was not identified. Further, despite the cluster-randomized design, there were significant imbalances in key demographic characteristics between groups at baseline, suggesting there may be unmeasured confounding, which detract from the ability to test the intervention effect. Generalizability is further limited in that participants within each work group could volunteer for the overall trial and further into the sub-study, assessing

accelerometer, suggesting these participants may not be representative of all employees working in similar jobs; however, this study was conducted in a diverse set of hospital work groups (e.g., food service, housekeeping, billing), enhancing the overall generalizability of the trial.

Conclusion

Overall, while we cannot conclude that the Working for You intervention improved PA among a subset of participants, the study contributes to our understanding of workplace interventions among diverse populations. Alternative strategies may be needed to promote health and address the significant structural barriers to numerous health behaviors among low income workers, a population disproportionately affected by health disparities.

Funding

This work was supported by the NIDDK; under Grants R01DK103760, 5P30DK020579, and P30DK092950. The content is solely the responsibility of the authors and does not necessarily represent the official view of the National Institutes of Health.

Corresponding should be addressed to

Rachel Tabak, PhD, RD

1 Brookings Drive, Campus Box 1196

St. Louis, MO 63130

Brown School at Washington University in St. Louis

rtabak@wustl.edu

314-935-0153



Jaime R. Strickland: 0000-0002-6296-7714



Maura Kepper: 0000-0002-5971-9189



Ann Marie Dale: 0000-0002-5624-8967



Bradley Evanoff: 0000-0003-0085-333X

Acknowledgements

Conflict of interest

No conflicts are declared by the authors.

Author contributions

Conceptualization, B. E., A. M. D., R. G. T., J. R. S.; Methodology, B. E., A. M. D., R. G. T., R. C., T. R.; Investigation, B. E., A. M. D., R. G. T., R. C., T. R.; Writing – Original Draft, R. G. T., Writing – Review & Editing, B. E., A. M. D., R. G. T., R. C., M. K., J. R. S., T.

R., Funding Acquisition, B. E., R. G. T.; Resources, B. E.;
Supervision, J. R. S.

Creative Commons License

This work is [licensed](#) under a [Creative Commons Attribution-Noncommercial 4.0 International License \(CC BY-NC 4.0\)](#).

Institutional approval

The study was approved by the Human Research Protection Office at Washington University in St. Louis (#201609029).

References

- Aneni, E. C., Roberson, L. L., Maziak, W., Agatston, A. S., Feldman, T., Rouseff, M., . . . Nasir, K. (2014). A systematic review of internet-based worksite wellness approaches for cardiovascular disease risk management: outcomes, challenges & opportunities. *PLoS One*, *9*(1), e83594. doi:10.1371/journal.pone.0083594
- Barbeau, E. M., Krieger, N., & Soobader, M. J. (2004). Working class matters: socioeconomic disadvantage, race/ethnicity, gender, and smoking in NHIS 2000. *Am J Public Health*, *94*(2), 269-278. doi:10.2105/ajph.94.2.269
- Blake, H., Suggs, L. S., Coman, E., Aguirre, L., & Batt, M. E. (2017). Active8! Technology-Based Intervention to Promote Physical Activity in Hospital Employees. *Am J Health Promot*, *31*(2), 109-118. doi:10.4278/ajhp.140415-QUAN-143
- Brown, S. A., Garcia, A. A., Zuniga, J. A., & Lewis, K. A. (2018). Effectiveness of workplace diabetes prevention programs: A systematic review of the evidence. *Patient Educ Couns*, *101*(6), 1036-1050. doi:10.1016/j.pec.2018.01.001
- Byers, T. E., Wolf, H. J., Bauer, K. R., Bolick-Aldrich, S., Chen, V. W., Finch, J. L., . . . Patterns of Care Study, G. (2008). The impact of socioeconomic status on survival after cancer in the United States : findings from the National Program of Cancer Registries Patterns of Care Study. *Cancer*, *113*(3), 582-591. doi:10.1002/cncr.23567
- Cairns, J. M., Bamba, C., Hillier-Brown, F. C., Moore, H. J., & Summerbell, C. D. (2015). Weighing up the evidence: a systematic review of the effectiveness of workplace interventions to tackle socio-economic inequalities in obesity. *J Public Health (Oxf)*, *37*(4), 659-670. doi:10.1093/pubmed/fdu077
- Clegg, L. X., Reichman, M. E., Miller, B. A., Hankey, B. F., Singh, G. K., Lin, Y. D., . . . Edwards, B. K. (2009). Impact of socioeconomic status on cancer incidence and stage at diagnosis: selected findings from the surveillance, epidemiology, and end results: National Longitudinal Mortality Study. *Cancer Causes Control*, *20*(4), 417-435. doi:10.1007/s10552-008-9256-0
- Coenen, P., Robroek, S. J. W., van der Beek, A. J., Boot, C. R. L., van Lenthe, F. J., Burdorf, A., & Oude Hengel, K. M. (2020). Socioeconomic inequalities in effectiveness of and compliance to workplace health promotion programs: an individual participant data (IPD) meta-analysis. *Int J Behav Nutr Phys Act*, *17*(1), 112. doi:10.1186/s12966-020-01002-w

- Craig, C. L., Marshall, A. L., Sjöström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., . . . Oja, P. (2003). International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc*, *35*(8), 1381-1395.
doi:10.1249/01.Mss.0000078924.61453.Fb
- Direito, A., Carraca, E., Rawstorn, J., Whittaker, R., & Maddison, R. (2017). mHealth Technologies to Influence Physical Activity and Sedentary Behaviors: Behavior Change Techniques, Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Ann Behav Med*, *51*(2), 226-239. doi:10.1007/s12160-016-9846-0
- Dowd, K. P., Szeklicki, R., Minetto, M. A., Murphy, M. H., Polito, A., Ghigo, E., . . . Donnelly, A. E. (2018). A systematic literature review of reviews on techniques for physical activity measurement in adults: a DEDIPAC study. *Int J Behav Nutr Phys Act*, *15*(1), 15. doi:10.1186/s12966-017-0636-2
- Folta, S. C., Paul, L., Nelson, M. E., Strogatz, D., Graham, M., Eldridge, G. D., . . . Seguin-Fowler, R. A. (2019). Changes in diet and physical activity resulting from the Strong Hearts, Healthy Communities randomized cardiovascular disease risk reduction multilevel intervention trial. *Int J Behav Nutr Phys Act*, *16*(1), 91. doi:10.1186/s12966-019-0852-z
- Goetzel, R. Z., & Ozminkowski, R. J. (2008). The health and cost benefits of work site health-promotion programs. *Annu Rev Public Health*, *29*, 303-323. doi:10.1146/annurev.publhealth.29.020907.090930
- Grosch, J. W., Alterman, T., Petersen, M. R., & Murphy, L. R. (1998). Worksite health promotion programs in the U.S.: factors associated with availability and participation. *Am J Health Promot*, *13*(1), 36-45. doi:10.4278/0890-1171-13.1.36
- Gu, M., Wang, Y., Shi, Y., Yu, J., Xu, J., Jia, Y., & Cheng, M. (2020). Impact of a group-based intervention program on physical activity and health-related outcomes in worksite settings. *BMC Public Health*, *20*(1), 935. doi:10.1186/s12889-020-09036-2
- Haskell, W. L., Lee, I. M., Pate, R. R., Powell, K. E., Blair, S. N., Franklin, B. A., . . . American Heart, A. (2007). Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation*, *116*(9), 1081-1093. doi:10.1161/CIRCULATIONAHA.107.185649
- Haughton, C. F., Silfee, V. J., Wang, M. L., Lopez-Cepero, A. C., Estabrook, D. P., Frisard, C., . . . Lemon, S. C. (2018). Racial/ethnic representation in lifestyle weight loss intervention studies in the United States: A systematic review. *Prev Med Rep*, *9*, 131-137. doi:10.1016/j.pmedr.2018.01.012
- Hawes, A. M., Smith, G. S., McGinty, E., Bell, C., Bower, K., LaVeist, T. A., . . . Thorpe, R. J., Jr. (2019). Disentangling Race, Poverty, and Place in Disparities in Physical Activity. *Int J Environ Res Public Health*, *16*(7).
doi:10.3390/ijerph16071193
- Howell, N. A., & Booth, G. L. (2022). The weight of place: Built environment correlates of obesity and diabetes. *Endocr Rev*.
doi:10.1210/endrev/bnac005

- Huang, Y., Hannon, P. A., Williams, B., & Harris, J. R. (2011). Workers' health risk behaviors by state, demographic characteristics, and health insurance status. *Prev Chronic Dis*, 8(1), A12. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/21159224>
- James, P., Hart, J. E., Hipp, J. A., Mitchell, J. A., Kerr, J., Hurvitz, P. M., . . . Laden, F. (2017). GPS-Based Exposure to Greenness and Walkability and Accelerometry-Based Physical Activity. *Cancer Epidemiol Biomarkers Prev*, 26(4), 525-532. doi:10.1158/1055-9965.EPI-16-0925
- Javed, Z., Haisum Maqsood, M., Yahya, T., Amin, Z., Acquah, I., Valero-Elizondo, J., . . . Nasir, K. (2022). Race, Racism, and Cardiovascular Health: Applying a Social Determinants of Health Framework to Racial/Ethnic Disparities in Cardiovascular Disease. *Circ Cardiovasc Qual Outcomes*, 15(1), e007917. doi:10.1161/CIRCOUTCOMES.121.007917
- Javed, Z., Maqsood, M. H., Amin, Z., & Nasir, K. (2022). Race and Ethnicity and Cardiometabolic Risk Profile: Disparities Across Income and Health Insurance in a National Sample of US Adults. *Journal of Public Health Management and Practice*, 28, S91-S100. doi:10.1097/phh.0000000000001441
- Jun, S. Y., Kim, J., Choi, H., Kim, J. S., Lim, S. H., Sul, B., & Hong, B. Y. (2019). Physical Activity of Workers in a Hospital. *Int J Environ Res Public Health*, 16(4). doi:10.3390/ijerph16040532
- Kelly, P., Fitzsimons, C., & Baker, G. (2016). Should we reframe how we think about physical activity and sedentary behaviour measurement? Validity and reliability reconsidered. *International Journal of Behavioral Nutrition and Physical Activity*, 13(1), 32.
- Kepper, M. M., Myers, C. A., Denstel, K. D., Hunter, R. F., Guan, W., & Broyles, S. T. (2019). The neighborhood social environment and physical activity: a systematic scoping review. *Int J Behav Nutr Phys Act*, 16(1), 124. doi:10.1186/s12966-019-0873-7
- Kogevinas, M., & Porta, M. (1997). Socioeconomic differences in cancer survival: a review of the evidence. *IARC Sci Publ*(138), 177-206. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/9353665>
- Kopicki, A., Van Horn, C., & Zukin, C. (2009). Healthy at work? unequal access to employer wellness programs. WorkTRENDS survey. vol 10. newark, new jersey: John J. Heldrich Center for Workforce Development, Edward J. Bloustein School of Planning and Public Policy, Rutgers University, 1.96-94.92.
- Kreuter, M. W., McQueen, A., Boyum, S., & Fu, Q. (2016). Unmet basic needs and health intervention effectiveness in low-income populations. *Prev Med*, 91, 70-75. doi:10.1016/j.ypmed.2016.08.006
- Kriska, A. M., Rockette-Wagner, B., Edelstein, S. L., Bray, G. A., Delahanty, L. M., Hoskin, M. A., . . . Group, D. R. (2020). The Impact of Physical Activity on the Prevention of Type 2 Diabetes: Evidence and Lessons Learned From the

- Diabetes Prevention Program, a Long-Standing Clinical Trial Incorporating Subjective and Objective Activity Measures. *Diabetes Care*, 44(1), 43-49.
- Lee, R. E., & Cubbin, C. (2009). Striding toward social justice: the ecologic milieu of physical activity. *Exerc Sport Sci Rev*, 37(1), 10-17. doi:10.1097/JES.0b013e318190eb2e
- Limb, E. S., Ahmad, S., Cook, D. G., Kerry, S. M., Ekelund, U., Whincup, P. H., . . . Harris, T. (2019). Measuring change in trials of physical activity interventions: a comparison of self-report questionnaire and accelerometry within the PACE-UP trial. *Int J Behav Nutr Phys Act*, 16(1), 10. doi:10.1186/s12966-018-0762-5
- Linnan, L. A., Sorensen, G., Colditz, G., Klar, D. N., & Emmons, K. M. (2001). Using theory to understand the multiple determinants of low participation in worksite health promotion programs. *Health Educ Behav*, 28(5), 591-607. Retrieved from <http://www.ncbi.nlm.nih.gov/htbin-post/Entrez/query?db=m&form=6&dopt=r&uid=11575688>
- Luo, J., Hendryx, M., & Wang, F. (2022). Mortality disparities between Black and White Americans mediated by income and health behaviors. *SSM Popul Health*, 17, 101019. doi:10.1016/j.ssmph.2021.101019
- Malik, S. H., Blake, H., & Suggs, L. S. (2014). A systematic review of workplace health promotion interventions for increasing physical activity. *Br J Health Psychol*, 19(1), 149-180. doi:10.1111/bjhp.12052
- Matthew, C. E. (2005). Calibration of accelerometer output for adults. *Medicine and Science in Sports and Exercise*, 37(11 Suppl), S512-522.
- Matthews, C. E., Chen, K. Y., Freedson, P. S., Buchowski, M. S., Beech, B. M., Pate, R. R., & Troiano, R. P. (2008). Amount of time spent in sedentary behaviors in the United States, 2003-2004. *Am J Epidemiol*, 167(7), 875-881. doi:10.1093/aje/kwm390
- Mazzucca, S., Arredondo, E. M., Hoelscher, D. M., Haire-Joshu, D., Tabak, R. G., Kumanyika, S. K., & Brownson, R. C. (2021). Expanding Implementation Research to Prevent Chronic Diseases in Community Settings. *Annu Rev Public Health*. doi:10.1146/annurev-publhealth-090419-102547
- McGowan, V. J., Buckner, S., Mead, R., McGill, E., Ronzi, S., Beyer, F., & Bamba, C. (2021). Examining the effectiveness of place-based interventions to improve public health and reduce health inequalities: an umbrella review. *BMC Public Health*, 21(1), 1888. doi:10.1186/s12889-021-11852-z
- McLaren, L. (2007). Socioeconomic status and obesity. *Epidemiol Rev*, 29, 29-48. doi:10.1093/epirev/mxm001
- NCI Division of Cancer Control and Population Sciences: Epidemiology and Genomics Research Program. (2020). SAS Programs for Analyzing NHANES 2003-2004 Accelerometer Data. Retrieved from <https://epi.grants.cancer.gov/nhanes-pam/>

- Parrish, A. T., Hammerback, K., Hannon, P. A., Mason, C., Wilkie, M. N., & Harris, J. R. (2018). Supporting the Health of Low Socioeconomic Status Employees: Qualitative Perspectives From Employees and Large Companies. *J Occup Environ Med, 60*(7), 577-583. doi:10.1097/JOM.0000000000001318
- Piercy, K. L., Troiano, R. P., Ballard, R. M., Carlson, S. A., Fulton, J. E., Galuska, D. A., . . . Olson, R. D. (2018). The Physical Activity Guidelines for Americans. *JAMA, 320*(19), 2020-2028. doi:10.1001/jama.2018.14854
- Prener, C. G., Gebauer, S., & Gilbert, K. L. (2021). Measuring the Effect of Place, Socioeconomic Status, and Racism on Coronary Heart Disease: Recent Trends and Missed Opportunities. *Current Epidemiology Reports, 8*(4), 190-199.
- Rongen, A., Robroek, S. J., van Lenthe, F. J., & Burdorf, A. (2013). Workplace health promotion: a meta-analysis of effectiveness. *Am J Prev Med, 44*(4), 406-415. doi:10.1016/j.amepre.2012.12.007
- Singer, J. D., Willett, J. B., & Willett, J. B. (2003). *Applied longitudinal data analysis: Modeling change and event occurrence*. Oxford: Oxford university press.
- Smith, M., Hosking, J., Woodward, A., Witten, K., MacMillan, A., Field, A., . . . Mackie, H. (2017). Systematic literature review of built environment effects on physical activity and active transport - an update and new findings on health equity. *Int J Behav Nutr Phys Act, 14*(1), 158. doi:10.1186/s12966-017-0613-9
- Song, Z., & Baicker, K. (2019). Effect of a Workplace Wellness Program on Employee Health and Economic Outcomes: A Randomized Clinical Trial. *JAMA, 321*(15), 1491-1501. doi:10.1001/jama.2019.3307
- Sorensen, G., McLellan, D. L., Sabbath, E. L., Dennerlein, J. T., Nagler, E. M., Hurtado, D. A., . . . Wagner, G. R. (2016). Integrating worksite health protection and health promotion: A conceptual model for intervention and research. *Prev Med, 91*, 188-196. doi:10.1016/j.ypmed.2016.08.005
- Stein, R. I., Strickland, J. R., Tabak, R. G., Dale, A. M., Colditz, G. A., & Evanoff, B. A. (2019). Design of a randomized trial testing a multi-level weight-control intervention to reduce obesity and related health conditions in low-income workers. *Contemp Clin Trials, 79*, 89-97. doi:10.1016/j.cct.2019.01.011
- Strickland, J. R., Kinghorn, A. M., Biver, S., Colvin, R., Dale, A. M., & Evanoff, B. A. (2020). *Tools for evaluating workplace participatory program implementation*. Paper presented at the APHA's 2020 VIRTUAL Annual Meeting and Expo (Oct. 24-28).
- Tabak, R. G., Strickland, J. R., Kirk, B., Colvin, R., Stein, R. I., Dart, H., . . . Evanoff, B. A. (2020). Pilot test of an interactive obesity treatment approach among employed adults in a university medical billing office. *Pilot Feasibility Stud, 6*, 57. doi:10.1186/s40814-020-00599-w

- Tabak, R. G., Strickland, J. R., Stein, R. I., Dart, H., Colditz, G. A., Kirk, B., . . . Evanoff, B. A. (2018). Development of a scalable weight loss intervention for low-income workers through adaptation of interactive obesity treatment approach (iOTA). *BMC Public Health, 18*(1), 1265. doi:10.1186/s12889-018-6176-0
- Terry, P. E. (2019). Workplace Health Promotion Is Growing Up but Confusion Remains About What Constitutes a Comprehensive Approach. *Am J Health Promot, 33*(6), 845-849. doi:10.1177/0890117119854618
- Thompson, T., McQueen, A., Croston, M., Luke, A., Caito, N., Quinn, K., . . . Kreuter, M. W. (2019). Social Needs and Health-Related Outcomes Among Medicaid Beneficiaries. *Health Educ Behav, 46*(3), 436-444. doi:10.1177/1090198118822724
- Troiano, R. P., Berrigan, D., Dodd, K. W., Masse, L. C., Tilert, T., & McDowell, M. (2008). Physical activity in the United States measured by accelerometer. *Medicine and Science in Sports and Exercise, 40*(1), 181-188. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18091006
- Tucker, J. M., Welk, G. J., & Beyler, N. K. (2011). Physical activity in U.S.: adults compliance with the Physical Activity Guidelines for Americans. *Am J Prev Med, 40*(4), 454-461. doi:10.1016/j.amepre.2010.12.016
- Tudor-Locke, C., Brashear, M. M., Johnson, W. D., & Katzmarzyk, P. T. (2010). Accelerometer profiles of physical activity and inactivity in normal weight, overweight, and obese U.S. men and women. *Int J Behav Nutr Phys Act, 7*, 60. doi:10.1186/1479-5868-7-60
- United States Department of Health and Human Services. (2020). Current Guidelines. *Office of Disease Prevention and Health Promotion, Office of the Assistant Secretary for Health, Office of the Secretary*. Retrieved from <https://health.gov/work/physical-activity/current-guidelines>
- Western, M. J., Armstrong, M. E. G., Islam, I., Morgan, K., Jones, U. F., & Kelson, M. J. (2021). The effectiveness of digital interventions for increasing physical activity in individuals of low socioeconomic status: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act, 18*(1), 148. doi:10.1186/s12966-021-01218-4
- Williams, W. M., Yore, M. M., & Whitt-Glover, M. C. (2018). Estimating physical activity trends among blacks in the United States through examination of four national surveys. *AIMS Public Health, 5*(2), 144-157. doi:10.3934/publichealth.2018.2.144
- Woods, L. M., Rachet, B., & Coleman, M. P. (2006). Origins of socio-economic inequalities in cancer survival: a review. *Ann Oncol, 17*(1), 5-19. doi:10.1093/annonc/mdj007
- Zhu, X., Yoshikawa, A., Qiu, L., Lu, Z., Lee, C., & Ory, M. (2020). Healthy workplaces, active employees: A systematic literature review on impacts of workplace environments on employees' physical activity and sedentary behavior. *Building and Environment, 168*, 106455.